

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/505, A61P 3/06	A1	(11) International Publication Number: WO 00/45819 (43) International Publication Date: 10 August 2000 (10.08.00)
(21) International Application Number: PCT/GB00/00285 (22) International Filing Date: 1 February 2000 (01.02.00) (30) Priority Data: 9902590.0 6 February 1999 (06.02.99) GB 9921062.7 8 September 1999 (08.09.99) GB (71) Applicant (for all designated States except US): AS-TRAZENECA UK LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB). (72) Inventor; and (75) Inventor/Applicant (for US only): RAZA, Ali [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). (74) Agent: BRYANT, Tracey; Global Intellectual Property, Patents, AstraZeneca UK Limited, Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: USE OF CHOLESTEROL-LOWERING AGENT (57) Abstract The invention concerns the use of particular oral dosages or dosage ranges of the compound (E) -7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof, to alter beneficially lipid levels or lipid ratios in a human patient in need thereof, as well as pharmaceutical compositions of said compound or salts adapted for oral administration which comprise such dosages, and methods of preparation thereof.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LJ	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

USE OF CHOLESTEROL-LOWERING AGENT

The present invention relates to the use of a cholesterol-lowering agent, and more particularly to the administration of a particular dose or dosage range of the HMG CoA reductase inhibitor, (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid and pharmaceutically acceptable salts thereof, hereinafter referred to as "the Agent" and illustrated (as the calcium salt) in formula I hereinafter. The invention further relates to the dosage range, start dose and dosage forms of the Agent.

10 The Agent is disclosed in European Patent Application, Publication No. 0521471, and in Bioorganic and Medicinal Chemistry, (1997), 5(2), 437-444 as an inhibitor of 3-hydroxy-3-methylglutaryl CoA reductase (HMG-CoA reductase) which is a major rate-limiting enzyme in cholesterol biosynthesis. The Agent is taught as useful in the treatment of hypercholesterolemia, hyperlipoproteinemia and atherosclerosis. HMG-CoA reductase inhibitors are the most widely used prescription medication for the treatment of
15 hypercholesterolaemia. A number of HMG-CoA reductase inhibitors are marketed, namely lovastatin, pravastatin, simvastatin, fluvastatin, atorvastatin and cerivastatin, and are collectively referred to as 'statins'. Despite the benefits of statin therapy, less than optimal results may be achieved in patients, due to the level of efficacy and safety achieved at the
20 recommended dosages of the currently marketed statins. Accordingly it is important to find dosages of alternative statins which beneficially alter lipid levels to a significantly greater extent than similar dosages of currently used statins and which have a similar or improved safety profile.

Surprisingly it has now been found that when dosed orally to patients with
25 hypercholesterolemia at particular dosages or in a particular dosage range the Agent lowers total cholesterol (TC) and low-density lipoprotein cholesterol (LDL-C) by an unexpected degree, and without any significant adverse side effects. When dosed at the same dosages or in the same dosage range, the Agent also modifies other lipoprotein levels (such as raising high density lipid cholesterol (HDL-C) levels, lowering triglyceride (TG) levels and lowering
30 apolipoprotein B-100 (Apo-B) levels) to an unexpected and beneficial extent, without any significant adverse side effects. Elevations of alanine aminotransferase (ALT) liver enzyme levels are reported for other HMG-CoA reductase inhibitors. Surprisingly it has now been

found that when the Agent is dosed at the dosages or in the dosage ranges discussed herein, clinically significant rises in these levels are less frequently observed.

Accordingly, one aspect of the present invention comprises a method of lowering LDL-C levels by 40% or more, and/or lowering total cholesterol levels by 30% or more,
5 and/or lowering triglyceride levels by 10% or more, and/or lowering apolipoprotein B-100 levels by 30% or more, and/or raising HDL-C levels by 5% or more, in a patient in need thereof, by administration of 5 to 80 mg per day of the Agent.

A further aspect of the present invention comprises a method of treatment of hypercholesterolemia in a patient in need thereof by administration of 5 to 80 mg per day of
10 the Agent.

A further aspect of the present invention comprises a method of lowering LDL-C levels by 40% or more in a patient in need thereof by administration of 5 to 80 mg per day of the Agent.

A further aspect of the present invention comprises a method of lowering LDL-C
15 levels by 45% or more in a patient in need thereof by administration of 10 to 80 mg per day of the Agent.

A further aspect of the present invention comprises a method of lowering LDL-C levels by 50% or more in a patient in need thereof by administration of 10 to 80 mg per day of the Agent.

20 A further aspect of the present invention comprises a method of lowering LDL-C levels by 55% or more in a patient in need thereof by administration of 20 to 80 mg per day of the Agent.

A further aspect of the present invention comprises a method of lowering LDL-C levels by 60% or more in a patient in need thereof by administration of 40 to 80 mg per day of
25 the Agent.

A preferred feature of the present invention comprises a method of treatment as described above wherein the LDL-C level of the patient prior to treatment with the Agent, when measured after 12 hours of fasting (water permitted) is 4.1 mmol/litre or more.

A further aspect of the present invention comprises a method of lowering total
30 cholesterol levels by 30% or more in a patient in need thereof by administration of 10 to 80 mg per day of the Agent.

A further aspect of the present invention comprises a method of lowering total cholesterol levels by 35% or more in a patient in need thereof by administration of 10 to 80 mg per day of the Agent.

A further aspect of the present invention comprises a method of lowering total
5 cholesterol levels by 40% or more in a patient in need thereof by administration of 40 to 80 mg per day of the Agent.

A further aspect of the present invention comprises a method of treating a hypercholesterolemic patient to bring said patient within recommended NCEP guidelines (or other recognised guidelines) by administration of 5 to 80 mg per day of the Agent.

10 Recommended NCEP guidelines refer to the National Cholesterol Education Program - Adult Treatment Panel (NCEP-ATP) II target LDL-C levels, which are incorporated herein by reference. A summary of these guidelines is given in JAMA, June 16, 1993-Vol 629, No. 23, pages 3015-3023, particularly Figure 1, page 3018-3019. NCEP guidelines recommend that patients with clinically evident coronary heart disease (CHD), such as documented
15 atherosclerosis, are titrated to LDL-C concentrations less than or equal to 100 mg/dl. Other recognised guidelines includes the European Atherosclerosis Society (EAS) guidelines, which recommend aggressive management of lipids in such patients with a target LDL-C of 115 mg/dl. Where a patient has no clinically evident CHD, but has 2 or more risk factors for such a disease, NCEP guidelines recommend patients are titrated to LDL-C concentrations less
20 than 130 mg/ml. Where a patient has no clinically evident CHD and one or no risk factors, the NCEP guideline target is less than 160 mg/dl.

A further aspect of the present invention comprises a method of treating hypertriglyceridemia by administration of 5 to 80 mg per day of the Agent.

A further aspect of the present invention comprises a method of raising high density
25 lipid (HDL) levels in a patient in need thereof by administration of 5 to 80 mg per day of the Agent.

A further aspect of the present invention comprises a method of raising or maintaining HDL levels whilst lowering LDL-C levels in a patient in need thereof by administration of 5 to 80 mg per day of the Agent. Preferred levels of lowering LDL-C levels, whilst maintaining
30 or raising HDL levels, include the percentage levels of reduction in LDL-C given in the above methods.

A further aspect of the present invention comprises a method of lowering Apolipoprotein B levels in a patient in need thereof by administration of 5 to 80 mg per day of the Agent.

A further aspect of the present invention comprises a method of treatment of mixed
5 hyperlipidemia in a patient in need thereof by administration of 5 to 80 mg per day of the Agent.

A further aspect of the present invention comprises a method of treatment of atherosclerosis in a patient in need thereof by administration of 5 to 80 mg per day of the Agent.

10 When a dose range of 5 to 80 mg per day, 10 to 80 mg per day, 20 to 80 mg per day or 40 to 80 mg per day is referred to herein, other particular dosage ranges which are further independent aspects of the invention include (as appropriate) 10 to 80 mg per day, 10 to 60 mg per day, 10 to 40 mg per day, 5 to 40 mg per day, 5 to 20 mg per day, 10 to 20 mg per day, 20 to 60 mg per day, 20 to 40 mg per day and 40 to 60 mg per day.

15 Accordingly, for the avoidance of doubt, particular independent aspects of the present invention comprise the following:-

(1) a method of lowering LDL-C levels in a patient in need thereof by

- (i) 40% or more by administration of 5 to 20 mg per day of the Agent;
- (ii) 45% or more by administration of 10-20 (preferably 10) mg per day of the Agent;
- 20 (iii) 50% or more by administration of 10-20 (preferably 10) mg per day of the Agent;
- (iv) 55% or more by administration of 20-40 mg per day of the Agent; or
- (v) 60% or more by administration of 40-80 mg per day of the Agent;

(2) a method of lowering total cholesterol levels in a patient in need thereof by

- 25 (i) 30% or more by administration of 5 to 10 mg per day of the Agent;
- (ii) 35% or more by administration of 10 to 20 mg per day of the Agent;
- (iii) 40% or more by administration of 20-40 mg per day of the Agent; or
- (iv) 45% or more by administration of 40-80 mg per day of the Agent;

30 (3) a method of raising HDL-C levels in a patient in need thereof by

- (i) 5% or more by administration of 5 to 80 mg per day (particularly 5 to 40 mg per day, more particularly 10 to 20 mg per day, especially 10 mg per day) of the Agent;

- (ii) 8% or more by administration of 5 to 20 mg per day of the Agent; or
- (iii) 10% or more by administration of 5 to 10 mg per day (especially 10 mg per day) of the Agent;

5 (4) a method of lowering apolipoprotein B-100 levels in a patient in need thereof by

- (i) 30% or more by administration of 5 to 80 mg per day (particularly 5 to 40 mg per day, more particularly 10 to 20 mg per day, especially 10 mg per day) of the Agent;
- (ii) 35% or more by administration of 5 to 80 mg per day (particularly 5 to 40 mg per day, more particularly 10 to 20 mg per day, especially 10 mg per day) of the Agent;
- 10 (iii) 40% or more by administration of 10 to 80 mg per day (particularly 5 to 40 mg per day, more particularly 10 to 20 mg per day, especially 10 mg per day) of the Agent;
- (iv) 45% or more by administration of 20 to 80 mg per day of the Agent; or
- (v) 50% or more by administration of 40 to 80 mg per day of the Agent;

15

(5) a method of lowering triglyceride levels levels in a patient in need thereof by

- (i) 10% or more by administration of 5 to 80 mg per day (particularly 5 to 40 mg per day, more particularly 10 to 20 mg per day, especially 10 mg per day) of the Agent;
- or
- 20 (ii) 20% or more by administration of 20 to 40 mg per day of the Agent;

Further independent aspects of the present invention comprise the following:-

- (1) a method of lowering LDL-C levels in a patient in need thereof by 30% or more by administration of 1 to 5 mg per day, more particularly 2.5 to 5 mg per day, of the Agent;
- 25 (2) a method of lowering total cholesterol levels in a patient in need thereof by 30% or more by administration of 2.5 to 5 mg per day of the Agent;
- (3) a method of raising HDL-C levels in a patient in need thereof by 5 % or more by administration of 2.5 to 5 mg per day of the Agent;
- (4) a method of lowering apolipoprotein B-100 levels in a patient in need thereof by
- 30 30% or more by administration of 2.5 to 5 mg per day of the Agent; or
- (5) a method of lowering triglyceride levels in a patient in need thereof by 10% or more by administration of 2.5 to 5 mg per day of the Agent.

Further aspects of the present invention comprise the following:-

- 5 (1) a method of lowering LDL-C levels by 30% or more and raising HDL-C levels by 5% (more particularly 8%) or more in a patient in need thereof by administration of 1 to 5 mg per day of the Agent;
- (2) a method of lowering LDL-C levels by 40% or more and raising HDL-C levels by 5% (more particularly 8%) or more in a patient in need thereof by administration of 2.5 to 10 mg per day of the Agent;
- 10 (3) a method of lowering LDL-C levels by 45% or more and raising HDL-C levels by 5% (more particularly 8%) or more in a patient in need thereof by administration of 5 to 20 mg per day of the Agent;
- (4) a method of lowering LDL-C levels by 50% or more and raising HDL-C levels by 5% (more particularly 8%) or more in a patient in need thereof by administration of 10 to 20 mg per day of the Agent;
- 15 (5) a method of lowering LDL-C levels by 55% or more and raising HDL-C levels by 5% (more particularly 8%) or more in a patient in need thereof by administration of 20 to 40 mg per day of the Agent; or
- (6) a method of lowering LDL-C levels by 60% or more and raising HDL-C levels by 5% (more particularly 8%) or more in a patient in need thereof by administration of 40 to 80 mg per day of the Agent.
- 20

Further aspects of the present invention comprise the following:-

- 25 (1) a method of lowering LDL-C levels by 30% or more and lowering apolipoprotein B-100 levels by 25% or more by administration of 1 to 5 mg per day of the Agent;
- (2) a method of lowering LDL-C levels by 45% or more and lowering apolipoprotein B-100 levels by 35% or more by administration of 5 to 20 mg per day (particularly 5 to 10 mg per day) of the Agent;
- 30 (3) a method of lowering LDL-C levels by 50% or more and lowering apolipoprotein B-100 levels by 40% or more by administration of 10 to 20 mg per day of the Agent;

- (4) a method of lowering LDL-C levels by 55% or more and lowering apolipoprotein B-100 levels by 45% or more by administration of 20 to 40 mg per day of the Agent;
or
(5) a method of lowering LDL-C levels by 60% or more and lowering apolipoprotein B-100 levels by 50% or more by administration of 40 to 80 mg per day of the Agent.

Further aspects of the present invention comprise the following:-

- (1) a method of lowering LDL-C levels by 30% or more and lowering triglyceride levels by 10% or more by administration of 1 to 5 mg per day of the Agent;
(2) a method of lowering LDL-C levels by 45% or more and lowering triglyceride levels by 10% or more by administration of 5 to 20 mg per day (particularly 5 to 10 mg per day) of the Agent;
(3) a method of lowering LDL-C levels by 50% or more and lowering triglyceride levels by 10% or more by administration of 10 to 20 mg per day of the Agent;
(4) a method of lowering LDL-C levels by 55% or more and lowering triglyceride levels by 10% (more especially 20%) or more by administration of 20 to 40 mg per day of the Agent; or
(5) a method of lowering LDL-C levels by 60% or more and lowering triglyceride levels by 10% (more especially 20%) or more by administration of 40 to 80 mg per day of the Agent.

Further aspects of the present invention comprise the following:-

- (1) A method of raising HDL-C levels by 5 % (particularly 8% and more particularly 10%) or more and lowering apolipoprotein B-100 levels by 25 % or more in a patient in need thereof by administration of 1 to 5 mg per day of the Agent; or
(2) A method of raising HDL-C levels by 5 % (particularly 8% and more particularly 10%) or more and lowering triglyceride levels by 10 % or more in a patient in need thereof by administration of 5 to 80 mg per day (particularly 5 to 40 mg per day, more particularly 10 to 20 mg per day, especially 10 mg per day) of the Agent.

Further aspects of the present invention comprise the following:-

- 5 (1) a method of lowering LDL-C levels by 30% or more, raising HDL-C levels by 5% (more particularly 8%) or more and lowering triglyceride levels by 10% or more in a patient in need thereof by administration of 1 to 5 mg per day of the Agent;
- (2) a method of lowering LDL-C levels by 40% or more, raising HDL-C levels by 5% (more particularly 8%) or more and lowering triglyceride levels by 10% or more in a patient in need thereof by administration of 2.5 to 10 mg per day of the Agent;
- 10 (3) a method of lowering LDL-C levels by 45% or more, raising HDL-C levels by 5% (more particularly 8%) or more and lowering triglyceride levels by 10% or more in a patient in need thereof by administration of 5 to 20 mg per day of the Agent;
- (4) a method of lowering LDL-C levels by 50% or more, raising HDL-C levels by 5% (more particularly 8%) or more and lowering triglyceride levels by 10% or more in a patient in need thereof by administration of 10 to 20 mg per day of the Agent;
- 15 (5) a method of lowering LDL-C levels by 55% or more, raising HDL-C levels by 5% (more particularly 8%) or more and lowering triglyceride levels by 10% or more in a patient in need thereof by administration of 20 to 40 mg per day of the Agent; or
- (6) a method of lowering LDL-C levels by 60% or more, raising HDL-C levels by 5% (more particularly 8%) or more and lowering triglyceride levels by 10% or more in a patient in need thereof by administration of 40 to 80 mg per day of the Agent.
- 20

Further aspects of the present invention comprise the following:-

- 25 (1) A method of raising HDL-C levels by 5 % (particularly 8% and more particularly 10%) or more, lowering apolipoprotein B-100 levels by 25 % or more and lowering triglyceride levels by 10% or more in a patient in need thereof by administration of 1 to 5 mg per day of the Agent; or
- (2) A method of raising HDL-C levels by 5 % (particularly 8% and more particularly 10%) or more, lowering apolipoprotein B-100 levels by 25 % or more and lowering triglyceride levels by 10 % or more in a patient in need thereof by administration of 5
- 30

to 80 mg per day (particularly 5 to 40 mg per day, more particularly 10 to 20 mg per day, especially 10 mg per day) of the Agent.

Further aspects of the present invention comprise the following:-

5

(1) a method of reducing the LDL-C/HDL-C lipid ratio to below 2.2 in a patient in need thereof by administration of 1 to 5 mg per day of the Agent;

(2) a method of reducing the LDL-C/HDL-C lipid ratio to below 2.0 in a patient in need thereof by administration of 5 to 10 mg per day of the Agent;

10 (3) a method of reducing the LDL-C/HDL-C lipid ratio to below 1.5 in a patient in need thereof by administration of 20 to 40 mg per day of the Agent;

(4) a method of reducing the LDL-C/HDL-C lipid ratio to below 1.2 in a patient in need thereof by administration of 80 mg per day of the Agent;

15 (5) a method of reducing the non-HDL-C/HDL-C lipid ratio to below 2.6 in a patient in need thereof by administration of 1 to 5 mg per day of the Agent;

(6) a method of reducing the non-HDL-C/HDL-C lipid ratio to below 2.2 in a patient in need thereof by administration of 5 to 10 mg per day of the Agent;

(7) a method of reducing the non-HDL-C/HDL-C lipid ratio to below 2.0 in a patient in need thereof by administration of 20 to 40 mg per day of the Agent;

20 (8) a method of reducing the non-HDL-C/HDL-C lipid ratio to below 1.5 in a patient in need thereof by administration of 1 to 5 mg per day of the Agent;

(9) a method of reducing the TC/HDL-C lipid ratio to below 3.6 in a patient in need thereof by administration of 1 to 5 mg per day of the Agent;

25 (10) a method of reducing the TC/HDL-C lipid ratio to below 3.3 in a patient in need thereof by administration of 5 to 10 mg per day of the Agent;

(11) a method of reducing the TC/HDL-C lipid ratio to below 3.0 in a patient in need thereof by administration of 20 to 40 mg per day of the Agent; or

(12) a method of reducing the TC/HDL-C lipid ratio to below 2.5 in a patient in need thereof by administration of 80 mg per day of the Agent.

30

A particularly suitable starting dose of the Agent in the methods referred herein is 5 to 10 mg per day, especially 10 mg per day. After initiation and/or upon titration of the Agent,

lipid levels may be analysed and the dosage adjusted accordingly. A further aspect of the invention is therefore a method as defined above when the Agent is administered at a starting dose of 5 or 10 mg per day, for example a method of lowering LDL-C levels in a patient in need thereof by 40% or more by administration of 5 or 10 mg per day of the Agent. A further
5 aspect of the present invention comprises a dosing regime comprising administration of a starting dose of 5 or 10 mg per day of the Agent to a patient in need thereof, such as a patient with an LDL-C level of 160 mg/dl or greater with no chronic heart disease (CHD) or peripheral vascular disease (PVD) and 1 or no risk factors, a patient with an LDL-C level of greater than 130 mg/dl with no CHD or PVD and with 2 or more risk factors, or a patient with
10 an LDL-C level of greater than 100 mg/dl with clinically evident CHD or PVD. Another particular group of patients who are beneficially treated by administration of a start dose of 5 or 10 mg per day of the Agent includes, for example, those patients whose LDL-C level is >4 mmol/litre and/or whose TC level is >5 mmol/litre and/or whose HDL-C level is < 1 mmol/litre and/or whose TG level is > 2mmol/litre. A starting dose of 5 or 10 mg per day of
15 the Agent unexpectedly has a superior efficacy and a comparable or better safety profile compared to the starting doses of other statins, and is therefore particularly advantageous.

In carrying out the methods of the invention, the Agent will be administered to a patient in the form of a pharmaceutical composition. A further aspect of the invention is therefore a pharmaceutical composition which comprises 5 to 80 mg of the Agent together
20 with a pharmaceutically acceptable excipient or diluent. Particular pharmaceutical compositions which themselves are further independent aspects of the invention comprise, for example, 5 mg, 10 mg, 20 mg, 40 mg and 80 mg of the Agent together with a pharmaceutically acceptable excipient or diluent. The pharmaceutical compositions will be in the form of a conventional dosage unit form, for example, tablets or capsules. Accordingly, a
25 further aspect of the invention comprises, a tablet or capsule containing the Agent in the amounts given above. The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Preferably the Agent is administered as a single dose once daily.

It will be appreciated that for each method of treatment referred to above, a further
30 aspect of the invention comprises the use of the Agent in the amount specified for the manufacture of a medicament for use in said methods of treatment.

Preferably the Agent is bis[(E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl)](3R,5S)-3,5-dihydroxyhept-6-enoic acid] calcium salt. Where herein a dose of (or a pharmaceutical composition comprising) 5 mg, 10 mg, 20 mg, 40 mg or 80 mg of the Agent is referred to, this includes a dose (or
 5 pharmaceutical composition comprising) of 5.2 mg, 10.4 mg, 20.8 mg, 41.6 mg and 83.2 mg respectively of the calcium salt of formula 1. Such doses (or compositions) are calculated to provide the equivalent of 5 mg, 10 mg, 20 mg, 40 mg and 80 mg respectively of the free acid form when dosed.

10 Pharmaceutical compositions

The following Example illustrates, but is not intended to limit, pharmaceutical dosage forms which are suitable for use in the invention as defined herein:

	Capsule	mg
15	The Agent	5.0
	Lactose	42.5
	Corn Starch	20.0
	Microcrystalline cellulose	32.0
	Pregelatinised starch	3.3
20	Synthetic Hydrotalcite	1.1
	Magnesium stearate	1.1

Capsules containing 1, 2.5 or 10 mg of the Agent may be obtained similarly using more or less lactose as appropriate to maintain a total fill weight of 105 mg.

25

Preferred such formulations are those in which the Agent is bis[(E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl)]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid] calcium salt.

To illustrate the invention, a randomised, dose response parallel-group study with
 30 bis[(E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]-pyrimidin-5-yl]]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid] calcium salt (hereinafter referred to as ZD4522) and atorvastatin (ATORV) in subjects with primary hypercholesterolaemia was carried out.

Primary objectives

The primary objective of this trial was to estimate the dose-response relationship between the dose of ZD4522 and the percentage reduction of LDL-C from the baseline value
5 with respect to placebo.

Secondary objectives

Secondary objectives of this trial included:
to estimate the effect of 10 and 80 mg doses of atorvastatin on LDL-C levels;
to estimate the effects of ZD4522 and atorvastatin on HDL-C, TG, TC, apolipoprotein
10 A-1, apolipoprotein Lp(a), apolipoprotein B-100 levels and LDL- C (by indirect method);
to assess the pharmacokinetics of oral doses of 1, 2.5, 5, 10, 20, and 40 mg ZD4522
(capsule formulations) over a 6 week treatment period; and
to assess the tolerability and safety of ZD4522 in comparison with placebo.

15 Trial Design

After a 6-week dietary run-in, subjects were randomised to either atorvastatin doses (10 or 80 mg), supplied open labelled, or to placebo or 1 of 6 ZD4522 doses (supplied blinded). Analysis of the blinded portion of the trial addressed the primary objective. The open atorvastatin groups were included to obtain additional data on the starting and high
20 doses, of a proven cholesterol-lowering agent in this patient population.

Trial Plan

A total of 10 visits were made to each clinic. Three of these took place during the placebo controlled dietary run-in period (Weeks -6, -2 and -1), five during the treatment
25 phase (Week 0 = randomisation visit, Weeks +1,+2, +4 and +6) and two follow up visits (Weeks +8 and +10).

Number of Subjects

The primary endpoint on which the sample size is based is on percentage reduction
30 from baseline in LDL-C (LDL cholesterol) values. A sample size of 9 in each group will have 90% power to detect a difference in means of 25% between 2 groups, assuming that the common standard deviation is 15%, using a 2 group t-test with a 0.05 two-sided significance

level. This has been increased to 12 subjects per group to adjust for multiple comparisons of groups against placebo while preserving a power of at least 90% (based on simulations). This sample size also leads to an estimate of the dose-response curve for percentage decline in LDL-C with a width of the confidence band less than 10% for most of the dose range.

5

Inclusion Criteria

For inclusion in the dietary run-in period, subjects had to fulfil all of the following criteria:

- (1) fasting LDL cholesterol (>4.14 but <6.21 mmol/L);
- 10 (2) fasting levels of triglycerides (<3.39 mmol/L);
- (3) male subjects, age 18 to 70 years. Female subjects age 50 to 70 years;
- (4) absence of menstruation for more than two years and follicle stimulating hormone (FSH), luteinising hormone (LH) and oestradiol within the post-menopausal range (using local laboratory ranges);
- 15 (5) body mass index ≤ 30 kg/m².

For inclusion in the randomised treatment phase of the trial, subjects had to fulfil all of the following criteria:

- (6) fasting LDL cholesterol >4.14 mmol/L but <5.69 mmol/L from measurements taken at weeks -2 and -1 before randomisation, with the lower value within 15% of the higher value;
- 20 (7) fasting TG <3.39 mmol/L at Weeks -6, -2 and -1, before randomisation;
- (8) a Food Record Rating Score (FRR) ≤ 15 at Weeks -2 and -1, before randomisation, to demonstrate compliance with the American Heart Association Step -1 diet.

Exclusion Criteria

- 25 Any of the following was regarded as a criterion for exclusion from the trial:

- (1) Subjects using cholesterol lowering drugs (this therapy must have been discontinued at least 4 weeks before the start of the dietary run-in period. Subjects taking probucol should have discontinued 12 months before inclusion in this study).
- (2) History of serious or hypersensitivity reactions to other HMG-CoA reductase
- 30 inhibitors.

- (3) Subjects with active arterial disease such as unstable angina, myocardial infarction, transient ischaemic attack (TIA), cerebro vascular accident (CVA) or coronary artery bypass graft (CABG) within 6 months or angioplasty within 3 months of study entry.
- (4) History of malignancy except subjects whose only malignancy has been basal or squamous cell skin carcinoma. Female subjects who have a history of cervical dysplasia unless 3 consecutive normal cervical smears have been recorded prior to entry.
- (5) Uncontrolled hypertension (diastolic blood pressure ≥ 95 mm Hg). Subjects who are taking thiazide diuretics and beta blockers will not be excluded provided they are maintained at a stable dose from 3 months before starting the study and throughout the study.
- 10 (6) Diabetes mellitus and/or other metabolic endocrine disease known to be associated with alterations in plasma lipid levels (uncontrolled hypothyroidism defined as a TSH > 1.5 times the ULN at week -6, subjects with nephrotic syndrome).
- (7) Known homozygous familial hypercholesterolaemia or known type III hyperlipoproteinaemia (familial dysbetalipoproteinaemia).
- 15 (8) Use of concomitant medications detailed as follows:

DISALLOWED MEDICATIONS

CLASS OF DRUG	GENERIC NAME
Antibiotics/ antifungals	Erythromycin Base Erythromycin Ethyl Succinate, Acetyl Sulfisoxazole Rifampicin Fluconazole Ketaconazole Itraconazole
Anti-epileptics/ antidepressants	Phenytoin Phenobarbitone Fluoxetine Carbamazepine
Acne treatment	Isotretinoin
Antiulcer drugs	Cimetidine Cisapride
Systemic Steroids	Triamcinolone Acetonide Triamcinolone Diacetate Betamethasone Sodium Phosphate Betamethasone Acetate Hydrocortisone Hydrocortisone Acetate Hydrocortisone Sodium Phosphate Hydrocortisone Sodium Succinate Cortisone Acetate Dexamethasone Dexamethasone Acetate Dexamethasone Sodium
Anti arrhythmic	Digoxin
Anticoagulant	Warfarin
Antihistamine	Astemizole Terfenadine
Systemic Steroids Systemic Steroids continued	Phosphate Prednisolone Methylprednisolone Methylprednisolone Acetate Methylprednisolone Sodium Succinate Prednisolone Tebutate Prednisolone Sodium Phosphate Methyltestosterone Fluoxymesterone any hormone replacement therapies (HRT)

Oral hypoglycaemic agents	Tolbutamide
Lipid Regulation	Niacin/Nicotinic Acid Probucol Psyllium Preparations Clofibrate Cholestyramine Colestipol Hydrochloride Gemfibrozil Lovastatin Pravastatin Sodium Simvastatin Fluvastatin
Immunosuppressants	Lymphocyte Immune Globulin Rho(D) Immune Globulin Azathiopane Sodium Muromonab-CD3 Cyclosporin

- (9) History of alcohol and/or drug abuse.
- (10) Subjects with active liver disease or hepatic dysfunction as defined by elevations in liver enzymes (ALT, AST, γ GT, ALP) or bilirubin \geq 1.5 times ULN at any time before randomisation at Visit 4.
- (11) Subjects with a serum creatinine $> 180 \mu\text{mol/L}$ prior to randomisation.
- (12) Participation in another study less than 3 months before enrolment in the Dietary Run-in period.
- (13) Serum CK > 3 times ULN at Weeks -6 and -1 (Visit 1 and 3).
- 10 (14) Clinically significant ophthalmic abnormalities, simple refractive errors will be allowed.
- (15) Subjects who have received randomised medication cannot be re-entered in this study.
- (16) Subjects who have serious or unstable medical or psychological conditions which in the opinion of the investigator, would compromise the subject's safety or successful
- 15 participation in the study.
- (17) Subjects who are receiving hormone replacement therapy (HRT).
- (18) Subjects who are receiving digoxin and/or coumarin anti-coagulants.

Trial Methods

Summary

The subjects were requested to take the study drug once daily, not less than 3 hours after the evening meal. At Visits 1, 2, 3, 4, 6, 7, 8, 9 and 10 the trial subjects reported to the clinic after fasting for twelve hours (water permitted) and blood was drawn to determine fasting lipids. Subjects provided a urine sample, and blood samples for haematology and clinical biochemistry evaluation. Where lipid and clinical biochemistry measurements fell at the same visit, only one sample was taken. At Visits 4, 6, 7 and 8 pharmacokinetic samples were collected. Weight, blood pressure and heart rate were measured and recorded at all visits. At Visits 1 and 10 an ECG was performed.

The analytical laboratory used is certified for standardisation of lipid analysis as specified by the Standardisation programme of the Centres for Disease Control and Prevention and the National Heart, Lung and Blood Institute.

15 Analysis

(a) Lipid

Concentrations of the following lipids were determined LDL-C, HDL-C, TG, TC, apo A-I, apo A-II, Lp(a), apo B, LDL-C (by Friedewald equation and β -quantification method).

20 (b) Biochemistry

Concentrations of the following were determined, AST, ALT, CK¹, γ GT, ALP and total bilirubin, sodium, potassium, calcium, chloride, phosphate, bicarbonate creatinine, fasting blood glucose and serum thyroxine (T₄) and thyroid stimulating hormone (TSH).

Unless surgically sterilised female subjects had post menopausal status confirmed by measurement of FSH, LH and oestradiol at Visit 1.

Haematology

Full blood count was performed and included the following; erythrocyte count, haemoglobin, haematocrit, leucocyte cell count, platelets, red cell distribution width, percentage differential leukocyte count (including percentage large unstained cells), mean corpuscular volume, mean corpuscular haemoglobin, mean corpuscular haemoglobin concentration. A clotting screen was performed at Visits 1 and 10 only.

Urinalysis

At Visits 1, 3, 4, 5, 6, 7, 8, 9 and 10 urinalysis was performed on 10 ml mid-stream urine samples and tested in the clinic for the following using labstix /dipstix: pH, glucose, blood, ketones, protein, and bilirubin.

5 Results

1	<u>Study Arm</u>	<u>Mean % Lowering of LDL-C at week 6(Friedewald method)</u>			
	Placebo	8			
	ZD4522 1 mg	36			
	ZD4522 2.5 mg	43			
10	ZD4522 5 mg	45			
	ZD4522 10 mg	52			
	ZD4522 20 mg	59			
	ZD4522 40 mg	63			
	Atorvastatin 10 mg	44			
15	Atorvastatin 80 mg	59			

2	<u>Study Arm</u>	<u>Lipid parameter (mean % change from baseline at week 6)</u>			
		TC	TG	HDL-C	Apo-B
	Placebo	-5.3	-1.7	+2.8	-2.8
20	ZD4522 1 mg	-25.3	-16.9	+8.8	-28.8
	ZD4522 2.5 mg	-30.8	-12.3	+8.8	-34.6
	ZD4522 5 mg	-32.9	-34.8	+13.2	-37.6
	ZD4522 10 mg	-36.7	-12.0	+13.7	-42.3
	ZD4522 20 mg	-41.9	-26.2	+9.2	-48.9
25	ZD4522 40 mg	-46.1	-25.5	+9.4	-54.6
	Atorvastatin 10 mg	-32.0	-13.6	+6.8	-36.4
	Atorvastatin 80 mg	-46.1	-30.7	-3.2	-50.2

Note: In 2, a positive value indicates a mean % increase of the lipid parameter level and a
 30 negative value indicates a mean % reduction of the lipid parameter level

3	<u>Study Arm</u>	<u>Lipid Ratios (mean value at week 6)</u>		
		non HDL-C/HDL-C	TC/HDL-C	LDL-C/HDL-C
	Placebo	4.3	5.3	3.8
5	ZD4522 1 mg	2.5	3.5	2.2
	ZD4522 2.5 mg	2.5	3.5	2.1
	ZD4522 5 mg	2.2	3.2	1.9
	ZD4522 10 mg	2.2	3.2	1.8
	ZD4522 20 mg	1.8	2.8	1.4
10	ZD4522 40 mg	1.5	2.5	1.3
	Atorvastatin 10 mg	2.5	3.5	2.1
	Atorvastatin 80 mg	2.1	3.1	1.7

4 Effects in each study arm on percentage change of total cholesterol from baseline are
 15 given in Fig.1. Dosing began at BASELINE after an initial dietary run in period of 6 weeks,
 as per the clinical protocol above. Dosing stopped at 6 weeks after BASELINE.


5 Effects in each study arm on percentage LDL cholesterol (Friedewald equation
 calculation) percentage change from baseline are given in Fig 2. Dosing began at BASELINE
 20 after an initial dietary run in period of 6 weeks, as per the clinical protocol above. Dosing
 stopped at 6 weeks after BASELINE.

6 Effects on each study arm on HDL cholesterol percentage change from baseline are
 given in Fig. 3. Dosing began at BASELINE after an initial dietary run in period of 6 weeks,
 25 as per the clinical protocol above. Dosing stopped at 6 weeks after BASELINE.

7 Effects on each study arm on Apolipoprotein B-100 percentage change from baseline
 are given in Fig. 4. Dosing began at BASELINE after an initial dietary run in period of 6
 weeks, as per the clinical protocol above. Dosing stopped at 6 weeks after BASELINE.

8. Effects on each study arm on triglyceride percentage change from baseline are given in Fig. 5. Dosing began at BASELINE after an initial dietary run in period of 6 weeks, as per the clinical protocol above. Dosing stopped at 6 weeks after BASELINE

It will be appreciated that alternative or additional trials, of shorter or longer duration, may be carried out with the Agent to demonstrate the present invention, such as other randomized double blind, optionally placebo controlled, trials with selected doses of the Agent and selected doses of commercially available statins (e.g. atorvastatin, pravastatin, simvastatin) as comparators in subjects with hypercholesterolaemia with or without a history of CHD or PVD or with risk factors therefor, measuring changes in LDL-C and other lipid or lipoprotein fractions and the safety of the Agent in respect of, for example, serum transaminase levels.



Formula I

CLAIMS

1. The use of the compound (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid, or a
5 pharmaceutically acceptable salt thereof, for the manufacture of an oral dosage form comprising 5 to 80 mg of the compound for use in lowering LDL-C levels by 40% or more, and/or lowering total cholesterol levels by 30% or more, and/or lowering triglyceride levels by 10% or more, and/or lowering apolipoprotein B-100 levels by 30% or more, and/or raising HDL-C levels by 5% or more in a human patient in need thereof, wherein the compound is
10 administered at 5 to 80 mg per day.
2. The use of the compound or a pharmaceutically acceptable salt thereof as claimed in claim 1 for the manufacture of an oral dosage form comprising 5 to 10 mg of the compound for use in lowering LDL-C levels by 40% or more, and/or lowering total cholesterol levels by
15 30% or more, and/or lowering triglyceride levels by 10% or more, and/or lowering apolipoprotein B-100 levels by 30% or more, and/or raising HDL-C levels by 5% or more in a human patient in need thereof, wherein the compound is administered at 5 to 10 mg per day.
3. The use of the compound or a pharmaceutically acceptable salt thereof as claimed in
20 claim 2 for the manufacture of an oral dosage form comprising 5 to 10 mg of the compound for use in lowering LDL-C levels by 45% or more in a human patient in need thereof.
4. The use of the compound or a pharmaceutically acceptable salt thereof as claimed in claim 2 for the manufacture of an oral dosage form comprising 5 to 10 mg of the compound
25 for use in lowering total cholesterol levels by 35% or more in a human patient in need thereof.
5. The use of the compound or a pharmaceutically acceptable salt thereof as claimed in claim 2 for the manufacture of an oral dosage form comprising 5 to 10 mg of the compound for use in lowering triglyceride levels by 10% or more in a human patient in need thereof.
30
6. The use of the compound or a pharmaceutically acceptable salt thereof as claimed in claim 2 for the manufacture of an oral dosage form comprising 5 to 10 mg of the compound

for use in lowering apolipoprotein B-100 levels by 35% or more in a human patient in need thereof.

7. The use of the compound or a pharmaceutically acceptable salt thereof as claimed in claim 2 for the manufacture of an oral dosage form comprising 5 to 10 mg of the compound for use in raising HDL-C levels by 8% or more in a human patient in need thereof.

8. The use as claimed in any of claims 1 to 7 wherein the oral dosage form is administered as a single dose once daily.

10

9. A pharmaceutical composition adapted for oral administration which comprises 5 to 80 mg of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable diluent or carrier.

15

10. A pharmaceutical composition adapted for oral administration which comprises 5 to 10 mg of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable diluent or carrier.

20

11. A pharmaceutical composition as claimed in claim 9 or 10 which comprises 5.2 to 10.4 mg of the calcium salt of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid, together with a pharmaceutically acceptable diluent or carrier.

25

12. A method of preparing a pharmaceutical composition as claimed in claim 9, 10 or 11 which comprises admixing the compound or a pharmaceutically acceptable salt thereof with a pharmaceutically acceptable diluent or carrier.

30 13. A method of lowering LDL-C levels by 40% or more, and/or lowering total cholesterol levels by 30% or more, and/or lowering triglyceride levels by 10% or more, and/or lowering apolipoprotein B-100 levels by 30% or more, and/or raising HDL-C levels by 5% or

more, in a human patient in need thereof, by oral administration of 5 to 80 mg per day of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof.

5 14. A method as claimed in claim 13 wherein 5 to 10 mg per day of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof is administered.

15 15. A method as claimed in claim 14 which is a method of lowering LDL-C levels by 45%
10 or more.

16. A method as claimed in claim 14 which is a method of lowering total cholesterol levels by 35% or more.

15 17. A method as claimed in claim 14 which is a method of lowering triglyceride levels by 10% or more.

18. A method as claimed in claim 14 which is a method of lowering apolipoprotein B-100 levels by 35% or more.

20


19. A method as claimed in claim 14 which is a method of raising HDL-C levels by 10% or more.


20. A method of reducing the LDL-C/HDL-C lipid ratio to below 2.0 in a patient in need
25 thereof by administration of 5 to 10 mg per day of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof.


21. A dosing regime comprising administration of a dose of 5 to 10 mg per day of (E)-7-
30 [4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]-(3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof to a patient having


- (i) an LDL-C level of 160mg/dl or greater with no chronic heart disease or peripheral vascular disease and one or no risk factors for such a disease;
- (ii) an LDL-C level of greater than 130 mg/dl with no chronic heart disease or peripheral vascular disease and two or more risk factors for such a disease; or
- 5 (iii) an LDL-C level of greater than 100 mg/dl with clinically evident chronic heart disease or peripheral vascular disease

Fig. 1.










5/5



INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/00285

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K31/505 A61P3/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 521 471 A (SHIONOGI SEIYAKU KABUSHIKI KAISHA) 7 January 1993 (1993-01-07) cited in the application the whole document page 4, line 25-28 examples 1,7 claims 1-9 — —/—	1-21

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

29 May 2000

Date of mailing of the international search report

15/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3018

Authorized officer

Economou, D

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/00285

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MASAMICHI WATANABE ET AL.: "Synthesis and biological activity of methanesulfonamide pyrimidine- and N-methanesulfonyl pyrrole-substituted 3,5-dihydroxy-6-heptenoates, a novel series of HMG-CoA reductase inhibitors" BIOORG. MED. CHEM., vol. 5, no. 2, 1997, pages 437-444, XP000909731 cited in the application the whole document	1-21
P, X	GRAUL, A. ET AL.: "ZD-4522; Hypolipidemic HMG-CoA reductase inhibitor" DRUGS FUTURE, vol. 24, no. 5, 1999, pages 511-513, XP000882032 SPAIN the whole document	1-21

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Jona! Application No

PCT/GB 00/00285

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0521471 A	07-01-1993	CA 2072945 A	02-01-1993
		HU 61531 A	28-01-1993
		JP 2648897 B	03-09-1997
		JP 5178841 A	20-07-1993
		KR 9605951 B	06-05-1996
		US 5260440 A	09-11-1993